

Newton Series Inversion Method

Victor Elgersma

07 June 2026

Contents

1 Introduction	1
2 Inverting $z = \ln(1 + x)$	2
3 Inverting $z = \sin^{-1}(x)$	10
Bibliography	16

1 Introduction

In Newton's *Analysis by Equations of an Infinite Number of Terms* (1669/1711), Newton calculated the inverse of the functions $\ln(1 + x)$ and $\sin^{-1}(x)$, culminating in the first appearance of the series for the sine and cosine in a European manuscript (Dunham, 1991, p. 18; Newton & Stewart, 1745, p. 336-338, §42-46)

2 Inverting $z = \ln(1 + x)$

He first demonstrates his series inversion method by inverting $z = \ln(1 + x)$ to get $x = e^z - 1$:

43. Thus if from the Area ABDC of the Hyperbola ($\frac{1}{1+x} = y$) given I wanted to investigate the Base AB, calling the Area z , I extract the Root of this Equation z (ABCD) $= x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4$, &c. neglecting those Terms in which x is of more Dimensions than z is desired in the Quotient.




Figure 1: (Newton & Stewart, 1745, p. 337)

It is no accident that he refers to x as a 'Quotient', since it appears to be the result of a process which closely resembles long-division, as we shall see.

So Newton starts with the power series:

$$\ln(1 + x) \equiv z(x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5 + \dots \quad (1)$$

His aim is to find an expression like:

$$x(z) = \alpha z + \beta z^2 + \gamma z^3 + \delta z^4 + \dots \quad (2)$$

Newton is aware that if he wants to find coefficients of the inverse power series, up to z^5 all he needs to do is invert infinite series of $z(x)$ truncated after the term x^5 :

As if I would have z to rise to five Dimensions only in the Quotient, I neglect all the Terms $-\frac{1}{6}x^6 + \frac{1}{7}x^7 - \frac{1}{8}x^8$, &c. and extract the Root of this only $\frac{1}{5}x^5 - \frac{1}{4}x^4 + \frac{1}{3}x^3 - \frac{1}{2}x^2 + x - z = 0$.

Figure 2: (Newton & Stewart, 1745, p. 337)

So, the infinite series:

$$z(x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5 + \dots \quad (3)$$

and the finite series:

$$z_5(x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5 \quad (4)$$

... will have a different inverse in general¹, but their inverses will be identical for the terms up to z^5 . Therefore, Newton can focus on inverting just the finite series $z_5(x)$.

Newton next rewrites Equation 4 and puts all terms on the LHS:

$$-z + x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5 = 0 \quad (5)$$

We see clearly from the expansion that:

$$z = x + O(x^2) + \dots \quad (6)$$

Therefore, the first-order approximation for z , which we call z_1 , is:

$$z_1 = x \quad (7)$$

The first-order approximation for x is given by inverting the first-order approximation for z . In other words,

$$x_1 = z \quad (8)$$

or

$$x = z + O(z^2) + \dots \quad (9)$$

Now, Newton rewrites Equation 5 by giving each power of x a separate row:

$$0 = \begin{cases} +\frac{1}{5}x^5 \\ -\frac{1}{4}x^4 \\ +\frac{1}{3}x^3 \\ -\frac{1}{2}x^2 \\ + x \\ - z \end{cases} \quad (10)$$

Now we make our substitution $x = z + p$, where p is taken to be $O(z^2)$

$$0 = \begin{cases} +\frac{1}{5}(z + p)^5 \\ -\frac{1}{4}(z + p)^4 \\ +\frac{1}{3}(z + p)^3 \\ -\frac{1}{2}(z + p)^2 \\ + z + p \\ - z \end{cases} \quad (11)$$

Then he expands Equation 11 selectively as follows:

¹For an interactive applet that shows the successive inverses $x_n(z)$ of the truncated power series $z_n(x)$ for $n = 1 \dots 12$ see <https://math.vjbe.net>

$$0 = \begin{cases} +\frac{1}{5}z^5 + z^4p + \dots & 5 - 5 = 0 \\ -\frac{1}{4}z^4 - z^3p + \frac{3}{2}z^2p^2 + \dots & 5 - 4 = 1 \\ +\frac{1}{3}z^3 + z^2p + zp^2 + \frac{1}{3}zp^3 + \dots & 5 - 3 = 2 \\ -\frac{1}{2}z^2 - zp - \frac{1}{2}p^2 + 0 & 5 - 2 = 3 \\ +z^1 + p \\ -z^1 \end{cases} \quad (12)$$

We have kept the first ignored terms in grey, and point out that they are all of a higher order in z than z^5 . Newton uses the following rule to determine which terms to keep:

Rule; That after the first Term resulting from each Quantity that is collateral to it, I add no more Terms upon the right Hand than the Index of the Dimension of that first Term wants Units of the Index of the greatest Dimension. As in this Example, where the greatest Dimension is 5, I neglect all the Terms after z^5 , I put one after z^4 , and two only after z^3 . When the Root (x) to be extracted,

The *Units of the Index of the greatest dimension* is 5. The *Index of the Dimension of the First term* is 5 for the top row, and decreases by one for each row. Therefore $5 - 5 = 0$ and *no* additional terms are added after the z^5 term. The same logic leads us to conclude that one term is added after the z^4 term, and two after the z^3 term.

This means that terms like z^4p or z^2p^2 , which are $\sim O(z^6)$, will be dropped. These terms are irrelevant because we are only seeking to find the coefficients up to z^5 .

We are thus left with the following expression:

$$0 = \begin{cases} +\frac{1}{5}z^5 \\ -\frac{1}{4}z^4 - z^3p \\ +\frac{1}{3}z^3 + z^2p + zp^2 \\ -\frac{1}{2}z^2 - zp - \frac{1}{2}p^2 \\ +z + p \\ -z \end{cases} \quad (13)$$

Or, in Newton's original text:

$$\begin{array}{r}
+ \frac{1}{5}z^5 \text{ \Ô} \\
- \frac{1}{4}z^4 - z^3p \text{ \Ô} \\
+ \frac{1}{3}z^3 + z^2p + zp^2 \text{ \Ô} \\
- \frac{1}{2}z^2 - zp - \frac{1}{2}p^2 \\
+ z + p \\
- z
\end{array}$$

Figure 3: (Newton & Stewart, 1745, p. 337)

We remember that we are looking for the lowest term in the power series expansion of p (where p is the correction to the current approximation $x = z$). Since we assume $p \sim O(z^2)$, we can ignore terms like p^2 . Collecting like terms:

$$0 = -\frac{1}{2}z^2 + \frac{1}{3}z^3 - \frac{1}{4}z^4 + \frac{1}{5}z^5 + p(1 - z + z^2 - z^3) \quad (14)$$

Rearranging:

$$p = \frac{+\frac{1}{2}z^2 - \frac{1}{3}z^3 + \frac{1}{4}z^4 - \frac{1}{5}z^5}{1 - z + z^2 - z^3} \quad (15)$$

Since we only seek the lowest-order coefficient of p , it is enough to expand $(1 - z + z^2 - z^3)^{-1} = (1 + O(z))$, so

$$p = \left(\frac{1}{2}z^2 - \frac{1}{3}z^3 + \frac{1}{4}z^4 - \frac{1}{5}z^5\right)(1 + O(z)) = \frac{1}{2}z^2 + O(z^3) \quad (16)$$

So we have now found our second order expansion of $x(z)$:

$$x_2(z) = z + \frac{z^2}{2} + O(z^3) \quad (17)$$

Newton's table up until this point looks like this:

$z + p = x$	$ \begin{array}{r} + \frac{1}{5}z^5 \\ - \frac{1}{4}z^4 \\ + \frac{1}{3}z^3 \\ - \frac{1}{2}z^2 \\ + z \\ - z \end{array} $	$ \begin{array}{r} + \frac{1}{5}z^5 \text{ } \mathcal{O}c. \\ - \frac{1}{4}z^4 - z^3p \text{ } \mathcal{O}c. \\ + \frac{1}{3}z^3 + z^2p + zp^2 \text{ } \mathcal{O}c. \\ - \frac{1}{2}z^2 - zp - \frac{1}{2}p^2 \\ + z + p \\ - z \end{array} $
$\frac{1}{2}z^2 + q = p$		

Figure 4: The steps to find the first and second terms in the inversion of $\ln(1+x)$

Now Newton returns to Equation 13, and rewrites it in order of decreasing dimensions of p and then z :

$$0 = \begin{cases} zp^2 \\ -\frac{1}{2}p^2 \\ -z^3p \\ +z^2p \\ -zp \\ +p \\ +\frac{1}{5}z^5 \\ -\frac{1}{4}z^4 \\ +\frac{1}{3}z^3 \\ -\frac{1}{2}z^2 \end{cases} \quad (18)$$

In Newton's original text, this looks like this:

$x + p = x$	$ \begin{array}{r} + \frac{1}{5}x^5 \\ - \frac{1}{4}x^4 \\ + \frac{1}{3}x^3 \\ - \frac{1}{2}x^2 \\ + x \\ - x \end{array} $	$ \begin{array}{r} + \frac{1}{5}z^5 \text{ \Ô.} \\ - \frac{1}{4}z^4 - z^3p \text{ \Ô.} \\ + \frac{1}{3}z^3 + z^2p + zp^2 \text{ \Ô.} \\ - \frac{1}{2}z^2 - zp - \frac{1}{2}p^2 \\ + z + p \\ - z \end{array} $
$\frac{1}{2}z^2 + q = p$	$ \begin{array}{r} + zp^2 \\ - \frac{1}{2}p^2 \\ - z^3p \\ + z^2p \\ - zp \\ + p \\ + \frac{1}{5}z^5 \\ - \frac{1}{4}z^4 \\ + \frac{1}{3}z^3 \\ - \frac{1}{2}z^2 \end{array} $	

Figure 5: (Newton & Stewart, 1745, p. 337)

Now, he substitutes $p = \frac{1}{2}z^2 + q$, where q is assumed to be of the form $q = O(z^3) + \dots$. He uses the same rules to determine which terms to keep

$$0 = \begin{cases} +z(\frac{1}{2}z^2 + q)^2 \\ -\frac{1}{2}(\frac{1}{2}z^2 + q)^2 \\ -z^3(\frac{1}{2}z^2 + q) \\ +z^2(\frac{1}{2}z^2 + q) \\ -z(\frac{1}{2}z^2 + q) \\ +(\frac{1}{2}z^2 + q) \\ +\frac{1}{5}z^5 \\ -\frac{1}{4}z^4 \\ +\frac{1}{3}z^3 \\ -\frac{1}{2}z^2 \end{cases} = \begin{cases} \frac{1}{4}z^5 + z^3q \\ -\frac{1}{8}z^4 - \frac{1}{2}z^2q \\ -\frac{1}{2}z^5 - z^3q \\ +\frac{1}{2}z^4 + z^2q \\ -\frac{1}{2}z^3 - qz \\ +\frac{1}{2}z^2 + q \\ +\frac{1}{5}z^5 \\ -\frac{1}{4}z^4 \\ +\frac{1}{3}z^3 \\ -\frac{1}{2}z^2 \end{cases} \quad \begin{matrix} 5 - 5 = 0 \\ 5 - 4 = 1 \\ 5 - 5 = 0 \\ 5 - 4 = 1 \\ 5 - 3 = 2 \end{matrix} \quad (19)$$

Since q only enters linearly in Equation 19, we have a simple closed-form solution for q in terms of z :

$$x = z + \frac{1}{2}z^2 + q \quad (20)$$

where q is found by solving for q in Equation 19:

$$q = \frac{\frac{1}{20}z^5 - \frac{1}{8}z^4 + \frac{1}{6}z^3}{1 - z + \frac{1}{2}z^2} \quad (21)$$

This can be found by polynomial long division, as Newton explains in §44.2:

2. When I see that p , q , or r , &c. in the last resulting Equation, is found of one Dimension only, I seek it's Value, that is to say the remaining Terms, which are still to be added to the Quotient, by means of Division; as you see done here.

Figure 6: (Newton & Stewart, 1745, p. 338)

$$1 - z + \frac{1}{2}z^2 \Big/ \frac{1}{6}z^3 - \frac{1}{8}z^4 + \frac{1}{20}z^5 \left(\frac{1}{6}z^3 + \frac{1}{24}z^4 + \frac{1}{120}z^5 \right)$$

Figure 7: (Newton & Stewart, 1745, p. 337)

We have recreated the long division below:

$$\begin{array}{r} 1 - z + \frac{1}{2}z^2 \Big/ \frac{1}{6}z^3 - \frac{1}{8}z^4 + \frac{1}{20}z^5 \Big\backslash \frac{1}{6}z^3 \\ \underline{\phantom{1 - z + \frac{1}{2}z^2} \frac{1}{6}z^3} \\ 0 \end{array}$$

Figure 8: Step 1: Dividing $\frac{1}{6}z^3$ by 1

$$\begin{array}{r} 1 - z + \frac{1}{2}z^2 \Big/ \frac{1}{6}z^3 - \frac{1}{8}z^4 + \frac{1}{20}z^5 \Big\backslash \frac{1}{6}z^3 \\ \underline{-\frac{1}{6}z^3 + \frac{1}{6}z^4 - \frac{1}{12}z^5} \\ 0 + \frac{1}{24}z^4 + \frac{1}{30}z^5 \end{array}$$

Figure 9: Step 2: Calculating the remainder

$$\begin{array}{r} 1 - z + \frac{1}{2}z^2 \Big/ \frac{1}{6}z^3 - \frac{1}{8}z^4 + \frac{1}{20}z^5 \Big\backslash \frac{1}{6}z^3 + \frac{1}{24}z^4 \\ \underline{-\frac{1}{6}z^3 + \frac{1}{6}z^4 - \frac{1}{12}z^5} \\ 0 + \frac{1}{24}z^4 - \frac{1}{30}z^5 \end{array}$$

Figure 10: Step 3: Dividing $\frac{1}{24}z^4$ by 1

$$\begin{array}{r} 1 - z + \frac{1}{2}z^2 \Big/ \frac{1}{6}z^3 - \frac{1}{8}z^4 + \frac{1}{20}z^5 \Big\backslash \frac{1}{6}z^3 + \frac{1}{24}z^4 \\ \underline{-\frac{1}{6}z^3 + \frac{1}{6}z^4 - \frac{1}{12}z^5} \\ 0 + \frac{1}{24}z^4 - \frac{1}{30}z^5 \\ \underline{-\frac{1}{24}z^4 + \frac{1}{24}z^5} \\ 0 + \frac{1}{120}z^5 \end{array}$$

Figure 11: Step 4: Calculating the remainder

$$\begin{array}{r} 1 - z + \frac{1}{2}z^2 \Big/ \frac{1}{6}z^3 - \frac{1}{8}z^4 + \frac{1}{20}z^5 \Big\backslash \frac{1}{6}z^3 + \frac{1}{24}z^4 + \frac{1}{120}z^5 \\ \underline{-\frac{1}{6}z^3 + \frac{1}{6}z^4 - \frac{1}{12}z^5} \\ 0 + \frac{1}{24}z^4 - \frac{1}{30}z^5 \\ \underline{-\frac{1}{24}z^4 + \frac{1}{24}z^5} \\ 0 + \frac{1}{120}z^5 \end{array}$$

Figure 12: Step 5: Dividing $\frac{1}{120}z^5$ by 1 to yield the final term

At this point we can stop, remembering that we sought the coefficients up to z^5 . We could, of course, continue the division beyond z^5 but these terms would be meaningless as the inverse of the infinite series $z(x)$ and the inverse of the finite series $z_5(x)$ cease to be identical after the z^5 term. Thus, the first 5 terms of the expansion of $e^z - 1$ is:

$$e^z - 1 \equiv x = z + \frac{1}{2}z^2 + \frac{1}{6}z^3 + \frac{1}{24}z^4 + \frac{1}{120}z^5 + \dots \quad (22)$$

If we wanted to find the next coefficient, we would have had to start with one more power of x in the series for z , namely:

$$z_6(x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5 - \frac{1}{6}x^6 \quad (23)$$

And use the same method.

This method is closely related to the modern method of undetermined coefficients, except the latter makes it more explicit that we are assuming an equation of the form $\sum_i^\infty C_i x^i$, whereas Newton's method leaves that implicit. We believe that the parallel with long division with decimal numbers would have made the explicit assumption of an equation of a certain form to be pedantic. After all, when dividing two decimal numbers, we do not start by *assuming* an answer of a certain form (with a decimal point and infinite digits after the decimal). We simply work it out. Similarly, Newton just works out the series without worrying about stating its form to begin with.

The completed table is shown below:

$x = z + \frac{1}{2}z^2 + \frac{1}{6}z^3 + \frac{1}{4}z^4 + \frac{1}{10}z^5 \text{ \Ô.}$		
$z + p = x$	$\begin{aligned} &+ \frac{1}{2}z^2 \\ &- \frac{1}{4}z^4 \\ &+ \frac{1}{3}z^3 \\ &- \frac{1}{2}z^2 \\ &+ z \\ &- z \end{aligned}$	$\begin{aligned} &+ \frac{1}{2}z^2 \text{ \Ô.} \\ &- \frac{1}{4}z^4 - z^3p \text{ \Ô.} \\ &+ \frac{1}{3}z^3 + z^2p + zp^2 \text{ \Ô.} \\ &- \frac{1}{2}z^2 - zp - \frac{1}{2}p^2 \\ &+ z + p \\ &- z \end{aligned}$
$\frac{1}{2}z^2 + q = p$	$\begin{aligned} &+ zp^2 \\ &- \frac{1}{2}p^2 \\ &- z^3p \\ &+ z^2p \\ &- zp \\ &+ p \\ &+ \frac{1}{2}z^5 \\ &- \frac{1}{4}z^4 \\ &+ \frac{1}{3}z^3 \\ &- \frac{1}{2}z^2 \end{aligned}$	$\begin{aligned} &+ \frac{1}{4}z^5 \text{ \Ô.} \\ &- \frac{1}{8}z^4 - \frac{1}{2}z^2q \\ &- \frac{1}{2}z^5 \text{ \Ô.} \\ &+ \frac{1}{2}z^4 + z^2q \\ &- \frac{1}{2}z^3 - zq \\ &+ \frac{1}{2}z^2 + q \\ &+ \frac{1}{5}z^5 \\ &- \frac{1}{4}z^4 \\ &+ \frac{1}{3}z^3 \\ &- \frac{1}{2}z^2 \end{aligned}$
$1 - z + \frac{1}{2}z^2 - \frac{1}{6}z^3 + \frac{1}{8}z^4 - \frac{1}{10}z^5 (\frac{1}{6}z^3 + \frac{1}{4}z^4 + \frac{1}{10}z^5)$		

Figure 13: Series inversion algorithm for $z = \ln(1 + x)$. Answer ($x = e^z - 1$)

3 Inverting $z = \sin^{-1}(x)$.

Now that we have learned Newton's method by inverting $\ln(1 + x)$, let's apply it to inverting $\sin^{-1}(x)$:

$$\sin^{-1}(x) \equiv z(x) = x + \frac{1}{6}x^3 + \frac{3}{40}x^5 + \frac{5}{112}x^7 + \frac{35}{1152}x^9 + \dots \quad (24)$$

This series differs from $\ln(1 + x)$ in that it contains only odd powers, and only positive coefficients. The coefficients themselves are more complicated, however. This time we want a solution up to the power of z^9 , so we must consider the truncated series:

$$z = x + \frac{1}{6}x^3 + \frac{3}{40}x^5 + \frac{5}{112}x^7 + \frac{35}{1152}x^9 \quad (25)$$

The first order series is:

$$z = x + O(x^3) + \dots \quad (26)$$

Therefore, the inverse to first order is:

$$x = z + O(z^3) + \dots \quad (27)$$

As before, the first order inverse is $x_1 = z$. We find the next order by substituting $x = z + p$ into Equation 25. We rearrange it first into decreasing powers of x

$$0 = \begin{cases} +\frac{35}{1152}x^9 \\ +\frac{5}{112}x^7 \\ +\frac{3}{40}x^5 \\ +\frac{1}{6}x^3 \\ +x \\ -z \end{cases} \quad (28)$$

Now we substitute the expansion $x = z + p$

$$0 = \begin{cases} +\frac{35}{1152}(z+p)^9 \\ +\frac{5}{112}(z+p)^7 \\ +\frac{3}{40}(z+p)^5 \\ +\frac{1}{6}(z+p)^3 \\ +x \\ -z \end{cases} \quad (29)$$

Again, we don't need to expand all the terms, only the ones that will be of a lower order than x^9 , which we have done below.

$$0 = \begin{cases} +\frac{35}{1152}(z^9 + 9z^8p) + \dots & \frac{9-9}{2} = 0 \\ +\frac{5}{112}(z^7 + 7z^6p + 28z^5p^2) + \dots & \frac{9-7}{2} = 1 \\ +\frac{3}{40}(z^5 + 5z^4p + 10z^3p^2 + 10z^2p^3) + \dots & \frac{9-5}{2} = 2 \\ +\frac{1}{6}(z^3 + 3z^2p + 3zp^2 + p^3) + \dots & \frac{9-3}{2} = 3 \\ +z + p \\ -z \end{cases} \quad (30)$$

Again, we have added the next (ignored) term in grey. This step relies on the assumption that the lowest term in p will be $O(z^3)$. Today we might point out that $\sin^{-1}(x)$, having only odd terms in its power series expansion, is an odd function, and the inverse of an odd function is also odd.

Newton explains how we must modify his rule about which terms to keep for the case of an even or odd power series such as this one:

after z^4 , and two only after z^3 . When the Root (x) to be extracted, is every where of even or odd Dimensions, let this be the Rule: That after the first Term, resulting from each Quantity which is collateral to it, you add no more Terms towards the Right Hand, than what the Index of the Dimension of that first Term, wants Pairs of Units of the Index of the highest Dimension; or no more than what it wants Ternaries of Units, when the Indexes of the Dimensions of x differ by three Units; and so in others.

The "Index of the Highest Dimension" is 9. The "Index of the dimension of the first term" is 9 for the first row and drops by 2 in each row. "wants Pairs of Units" means we have to divide the difference by two to get the number of additional units on the right hand side.

To summarize, ignoring orders higher than $O(z^9)$ allows us to consider the (much simplified) Equation 31 rather than the full Equation 29

$$0 = \begin{cases} +\frac{35}{1152}z^9 \\ +\frac{5}{112}(z^7 + 7z^6p) \\ +\frac{3}{40}(z^5 + 5z^4p + 10z^3p^2) \\ +\frac{1}{6}(z^3 + 3z^2p + 3zp^2 + p^3) \\ +z + p \\ -z \end{cases} \quad (31)$$

We now want to find the coefficient of the z^3 assumed to be the leading coefficient of p , so we take Equation 30, drop all higher powers of p , and solve for p :

$$0 = p \left(\frac{35}{112}z^6 + \frac{15}{40}z^4 + \frac{3}{6}z^2 + 1 \right) + \frac{1}{6}z^3 + \frac{3}{40}z^5 + \frac{5}{112}z^7 + \frac{35}{1152}z^9 \quad (32)$$

$$p = \frac{-\frac{1}{6}z^3 - \frac{3}{40}z^5 - \frac{5}{112}z^7 - \frac{35}{1152}z^9}{1 + \frac{1}{2}z^2 + \frac{15}{40}z^4 + \frac{35}{112}z^6} \quad (33)$$

$$p = -\frac{1}{6}z^3 + O(z^5) \quad (34)$$

We have thus found the third-order approximation:

$$x_3(z) = z - \frac{1}{6}z^3 \quad (35)$$

And thus we know the first two terms of our infinite power series expansion:

$$x(z) = z - \frac{1}{6}z^3 + \dots \quad (36)$$

We now rewrite Equation 30 in terms of decreasing powers of p and z , just like we did before (Equation 18) when calculating the inverse of $\ln(1+x)$ in Section 2. This arrangement makes it easier to substitute successive approximations for p , since terms of the same degree in p are grouped together.

$$0 = \begin{cases} +\frac{1}{6}p^3 \\ +\frac{3}{4}z^3p^2 \\ +\frac{1}{2}zp^2 \\ +\frac{35}{112}z^6p \\ +\frac{15}{40}z^4p \\ +\frac{1}{2}z^2p \\ p \\ +\frac{35}{1152}z^9 \\ +\frac{5}{112}z^7 \\ +\frac{3}{40}z^5 \\ +\frac{1}{6}z^3 \\ +z \\ -z \end{cases} \quad (37)$$

We now use the substitution $p = -\frac{1}{6}z^3 + q$ into Equation 37 where we assume that the leading coefficient of q is $O(z^5)$. Again, we use Newton's rule to discard terms of higher order than z^9 :

$$0 = \begin{cases} +\frac{1}{6}\left(-\frac{1}{6^3}z^9\right) + \dots & \frac{9-9}{2} = 0 \\ +\frac{3}{4}z^3\left(\frac{1}{36}z^6\right) + \dots & \frac{9-9}{2} = 0 \\ +\frac{1}{2}z\left(\frac{1}{36}z^6 \boxed{-\frac{1}{3}qz^3} + \dots\right) & \frac{9-7}{2} = \boxed{1} \\ +\frac{35}{112}z^6\left(-\frac{1}{6}z^3\right) + \dots & \frac{9-9}{2} = 0 \\ +\frac{15}{40}z^4\left(-\frac{1}{6}z^3 + \boxed{q}\right) & \frac{9-7}{2} = \boxed{1} \\ +\frac{1}{2}z^2\left(-\frac{1}{6}z^3 + q\right) & \frac{9-5}{2} = 2 \\ -\frac{1}{6}z^3 + q \\ +\frac{35}{1152}z^9 \\ +\frac{5}{112}z^7 \\ +\frac{3}{40}z^5 \\ +\frac{1}{6}z^3 \\ +z \\ -z \end{cases} \quad (38)$$

We can expand the brackets and add like terms:

$$0 = \begin{cases} \left(-\frac{1}{1296} + \frac{1}{48} - \frac{35}{672} + \frac{35}{1152}\right)z^9 \\ \left(+\frac{1}{72} - \frac{15}{240} + \frac{5}{112}\right)z^7 \\ \left(+\frac{15}{40} - \frac{1}{6}\right)qz^4 \\ \left(-\frac{1}{12} + \frac{3}{40}\right)z^5 \\ +\frac{1}{2}z^2q \\ -\frac{1}{6}z^3 + q \\ +\frac{1}{6}z^3 \end{cases} \quad (39)$$

At this point we find that our equation is linear in q , which means we can solve it using polynomial division:

$$0 = \begin{cases} -\frac{17}{10368}z^9 \\ -\frac{1}{252}z^7 \\ +\frac{5}{24}qz^4 \\ -\frac{1}{120}z^5 \\ +\frac{1}{2}z^2q \\ q \end{cases} \quad (40)$$

$$\frac{1}{120}z^5 + \frac{1}{252}z^7 + \frac{17}{10368}z^9 = q\left(1 + \frac{1}{2}z^2 + \frac{5}{24}z^4\right) \quad (41)$$

$$\frac{\frac{1}{120}z^5 + \frac{1}{252}z^7 + \frac{17}{10368}z^9}{1 + \frac{1}{2}z^2 + \frac{5}{24}z^4} = q \quad (42)$$

We can calculate the remaining terms by straightforward polynomial division:

Figure 14: Step 1: Dividing $\frac{1}{120}z^5$ by 1

Figure 15: Step 2: Calculating the remainder

Figure 16: Step 3: Dividing $-\frac{1}{5040}z^7$ by 1

Figure 17: Step 4: Calculating the remainder

Figure 18: Step 4: Dividing $\frac{1}{362880}z^9$ into 1

We found that

$$q = \frac{1}{120}z^5 - \frac{1}{5040}z^7 + \frac{1}{362880}z^9 \quad (43)$$

And therefore

$$x = z - \frac{1}{6}z^3 + q \quad (44)$$

$$x = z - \frac{1}{6}z^3 + \frac{1}{120}z^5 - \frac{1}{5040}z^7 + \frac{1}{362880}z^9 + O(z^{11}) \quad (45)$$

We recognize this as the first few terms of the formula:

$$x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} z^{2n+1} \quad (46)$$

45. If from the Arch αD given the Sine AB was required; I extract the Root of the Equation found above, viz. $x = z - \frac{1}{6}z^3 + \frac{1}{120}z^5 - \frac{1}{5040}z^7 + \frac{1}{362880}z^9 + O(z^{11})$ (it being supposed that $AB = x$, $\alpha D = z$, and $A\alpha = 1$) by which I find $x = z - \frac{1}{6}z^3 + \frac{1}{120}z^5 - \frac{1}{5040}z^7 + \frac{1}{362880}z^9 \text{ \&c.}$

Figure 19: (Newton & Stewart, 1745, p. 338)

To calculate the z^{13} we would have to start with the expansion of $z(x)$ up to order x^{15}

Newton does not give us a table for his inversion of $\sin^{-1}(x)$ as he did for $\ln(1+x)$. But if he had, we can be fairly confident it would have looked something like this:

$x = z - \frac{1}{6}z^3 + \frac{1}{120}z^5 - \frac{1}{5040}z^7 + \frac{1}{362880}z^9 + O(z^{11})$		
$x = z + p$	$ \begin{aligned} &+ \frac{35}{1152}x^9 \\ &+ \frac{5}{112}x^7 \\ &+ \frac{3}{40}x^5 \\ &+ \frac{1}{6}x^3 \\ &+ x \\ &- z \end{aligned} $	$ \begin{aligned} &+ \frac{35}{1152}z^9 \\ &+ \frac{5}{112}(z^7 + 7z^6p) \\ &+ \frac{3}{40}(z^5 + 5z^4p + 10z^3p^2) \\ &+ \frac{1}{6}(z^3 + 3z^2p + 3zp^2 + p^3) \\ &+ z + p \\ &- z \end{aligned} $
$p = -\frac{1}{6}z^3 + q$	$ \begin{aligned} &+ \frac{1}{6}p^3 \\ &+ \frac{3}{4}z^3p^2 \\ &+ \frac{1}{2}zp^2 \\ &+ \frac{35}{112}z^6p \\ &+ \frac{15}{40}z^4p \\ &+ \frac{1}{2}z^2p \\ &p \\ &+ \frac{35}{1152}z^9 \\ &+ \frac{5}{112}z^7 \\ &+ \frac{3}{40}z^5 \\ &+ \frac{1}{6}z^3 \\ &+ z \\ &- z \end{aligned} $	$ \begin{aligned} &+ \frac{1}{6}\left(-\frac{1}{6^3}z^9\right) + \dots \\ &+ \frac{3}{4}z^3\left(\frac{1}{36}z^6\right) + \dots \\ &+ \frac{1}{2}z\left(\frac{1}{36}z^6 - \frac{1}{3}qz^3 + \dots\right) \\ &+ \frac{35}{112}z^6\left(-\frac{1}{6}z^3\right) + \dots \\ &+ \frac{15}{40}z^4\left(-\frac{1}{6}z^3 + q\right) \\ &+ \frac{1}{2}z^2\left(-\frac{1}{6}z^3 + q\right) \\ &-\frac{1}{6}z^3 + q \\ &+ \frac{35}{1152}z^9 \\ &+ \frac{5}{112}z^7 \\ &+ \frac{3}{40}z^5 \\ &+ \frac{1}{6}z^3 \\ &+ z \end{aligned} $
$ \left(1 + \frac{1}{2}z^2 + \frac{5}{24}z^4\right) \left(\frac{1}{120}z^5 + \frac{1}{252}z^7 + \frac{17}{10368}z^9\right) = \frac{1}{120}z^5 - \frac{1}{5040}z^7 + \frac{1}{362880}z^9 $		

Compare with Figure 13.

Bibliography

Dunham, W. (1991). *Journey through Genius: The Great Theorems of Mathematics*. Penguin Books.

Newton, I., & Stewart, J. (1745). *Sir Isaac Newton's Two Treatises of the Quadrature of Curves, and Analysis by Equations of an Infinite Number of Terms Explained*. Printed by James Bettenham for J. Nourse. https://books.google.it/books?id=noQ_AAAAcAAJ